UCRL- 94778
PREPRINT

INVERSE COMPTON MODEL OF GAMMA RAY BURST SPECTRA

CIRCULATION COPY
SUBJECT TO RECALL
IN TWO WEEKS

W. M. Howard and E. P. Liang
University of California
Lawrence Livermore National Laboratory
Livermore, CA 94550

This paper was prepared for submittal to the
International Astronomical Union
Symposium No. 125
Nanjing, China
May 25-30, 1986

June 1986

Lawrence Livermore National Laboratory

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement recommendation, or favoring of the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

INVERSE COMPTON MODEL OF GAMMA RAY BURST SPECTRA*

W. M. Howard and E. P. Liang
Physics Department, LLNL
University of California
P.O. Box 808
Livermore, CA 94550 U.S.A.

ABSTRACT. We study gamma ray spectra produced by the inverse Compton upscattering of soft photons by relativistic electrons with a one dimensional momentum distribution, which is relevant to gamma ray burst if the source magnetic field is strong enough so that the synchrotron cooling time of transverse energy becomes much shorter than isotropization time via couloub or Compton collisions. We find that for high electron longitudinal temperatures the output power is strongly beamed in the momentum direction and the spectrum softens rapidly with increasing view angle from the momentum direction.

Even if the gamma rays of gamma bursters originate from regions of magnetic fields as low as 10^{10} - 10^{11} G, the synchrotron cooling time of transverse momentum ($\sim 10^{-14}$ s$(10^{11}$ G/B$)^2$) may still be much shorter than the isotropization time via coulomb or Compton collisions. In that case the electron momentum distribution will be essentially 1-D along the field lines. The hardest photons will be produced by inverse Compton upscattering of synchrotron and other soft photons by these 1-D relativistic electrons. We have computed the inverse Compton spectra with Monte Carlo techniques for Maxwellian electron distributions, varying the scattering depth, longitudinal temperature and input soft photon spectrum. The major findings include:

a) upscattered power is strongly beamed along momentum direction;
b) beam narrows with increasing electron temperature, with half angle less than 2° for temperatures above 1 MeV;
c) for a given temperature, beam narrowest near unit scattering depth;
d) spectrum softens rapidly for increasing view angle away from momentum direction. For further details refer to Ref. 1.

*Work performed under the auspices of the U.S. DOE by the LLNL under contract number W-7405-ENG-48.

References

1. W. M. Howard, E. P. Liang and E. Canfield, Astrophys. J. submitted (1986).